

Order Form

AREA RESERVED FOR PHARMACY



Prescriber Information

Practice Name

Street Address

City

State

ZIP

Phone

Fax

Patient Information

Name

DOB

Street Address

City

State

ZIP

Phone

Email

Allergies

Sex

Pharmacy To Dispense

Sildenafil (choose Sig:)

☐ 25 mg (1 oral tablet, 1 hour prior to intimacy)☐ 50 mg (1 oral tablet, 1 hour prior to intimacy)☐ 100 mg (1 oral tablet, 1 hour prior to intimacy)Quantity ☐ 30 tab ☐ 60 tab ☐ 90 tab Other:____ Refill:____

Tadalafil (choose Sig:)

☐ 2.5 mg (1 oral tablet, 1 hour prior to intimacy)☐ 5 mg (1 oral tablet, 1 hour prior to intimacy)☐ 10 mg (1 oral tablet, 1 hour prior to intimacy)☐ 20 mg (1 oral tablet, 1 hour prior to intimacy)Quantity ☐ 30 tab ☐ 60 tab ☐ 90 tab Other:____ Refill:____

Price Reference for Men's Health Oral ED Meds

Sildenafil: 25mg - \$1/tab -- 50mg - \$1.50/tab -- 100mg - \$2/tab

Tadalafil: 2.5mg - \$1/tab -- 5mg - \$1.50/tab -- 10mg - \$2/tab -- 20mg - \$2.50/tab

* Men's Health Oral ED Tabs must be dispensed in increments of 30 to receive full discount. *

Quantity ☐ 30 tab ☐ 60 tab ☐ 90 tab Other:____ Refill:____☐ **Bi-Mix** Papavarine 30 mg/mL and Phentolamine 2 mg/mL (choose Sig:)☐ **Tri-Mix 1** Papavarine 30 mg/mL and Phentolamine 1 mg/mL, and Alprostadil 10 mcg/mL (choose Sig:)☐ **Tri-Mix 2** Papavarine 30 mg/mL and Phentolamine 2 mg/mL, and Alprostadil 20 mcg/mL (choose Sig:)☐ 2.5 mL (1 injection, 15-50 units, 5-20 minutes prior to intimacy)☐ 5 mL (1 injection, 15-50 units, 5-20 minutes prior to intimacy)☐ 7.5 mL (1 injection, 15-50 units, 5-20 minutes prior to intimacy)☐ 10 mL (1 injection, 15-50 units, 5-20 minutes prior to intimacy)Quantity ☐ 1 vial ☐ 2 vials ☐ 3 vials Other:____ Refill:____

Directions for Use

Tablets

Take 1 tablet by mouth, with or without food, 1 hour prior to sexual relations. Use caution when taking with other blood pressure lowering medications, including nitroglycerin.

Injections

Administered as a penile self-injection. A maximum of three injections per week, but you must wait 24 hours before taking another dose. And make sure you alternate between sides of the penis for each injection.

Special Instructions

Prescriber Section

I certify that the above patient has been informed of the risks of using these drugs in combination with other blood pressure lowering medications, including nitroglycerin.

Prescriber Name

Supervising Physician

NPI

Date/Time

Prescriber Signature

Send the completed form to Lake Country Compounding & Wellness

Fax: 762-777-7002