

Commonly Requested Wound Care Compounding Ideas

Treatment of Wounds/Burns Without Infection

- Aloe Vera/Vitamin E/Allantoin/Glycosaminoglycans Spira-Wash™ Gel
- Allantoin 2%/Vitamin D3 1000 IU/Gm/Aloe Vera 0.5% Spira-Wash™ Gel
- Silver Sulfadiazine 1% Spira-Wash™ Gel
- Collagenase 250 U/Gm Spira-Wash™ Gel
- Phenytoin 5%/Misoprostol 0.0024%/Arginine HCl 1% Spira-Wash™ Gel
- Phenytoin 5%/Misoprostol 0.0024%/Arginine HCl 1%/Sodium Hyaluronate 0.5% Spira-Wash™ Gel
- Mupirocin 5% Topical Gel (Spira-Wash™)
- Ketoprofen 2%/Lidocaine 2%/Misoprostol 0.0024%/Phenytoin 2%/Aloe Vera 0.2% Spira-Wash™ Gel
- Phenytoin 5%/Misoprostol 0.0024% Spira-Wash™ Gel
- Phenytoin 5%/Misoprostol 0.0024%/Nifedipine 2% Spira-Wash™ Gel
- Phenytoin Sodium 2%/Nifedipine 5%/Tranilast 1%/Misoprostol 0.0024% Topical Gel (PracaSil™-Plus) Combines Spirawash and Pracasil together.
- Phenytoin 5%/Misoprostol 0.0024% Topical Gel
- Phenytoin 2%/Misoprostol 0.0024%/Lidocaine 2%/Bupivacaine HCl 0.2%/Diphenhydramine HCl 1%/Aloe Vera 0.2% Polyox Bandage Polyox bandage is a good base for oozing wounds. Ask the pharmacist for Patient Instructions for Use of External Wound Bandage.
- Ketoprofen 2%/Lidocaine 2%/Misoprostol 0.0024%/Phenytoin 2%/Aloe Vera 0.2% Topical Cream
- Aloe Vera/Vitamin E/Collagen/Allantoin/Lanolin Topical Cream



Treatment of Infected Wounds/Burns

- Metronidazole 4%/Hydrocortisone 1%/Zinc Sulfate 2%/Silver Sulfadiazine 1% Spira-Wash™ Gel
- Levofloxacin 2%/Mupirocin 4%/Itraconazole 1% Spira-Wash™ Gel

These Are Just a Few Ideas. Contact Us to Find Out More, Today!



Katherine Hunt, PharmD, RPh
Tiffany Wright, PharmD, RPh
1101 W. Main St., Suite F • League City, Texas 77573
Ph: (832) 905-2961 • Fax: (832) 905-3142
www.gccompoundingrx.com

Commonly Requested Wound Care Compounding Ideas

- Misoprostol 0.0024%/Ciprofloxacin HCl 2% Spira-Wash™ Gel
- Misoprostol 0.0024%/Metronidazole 2%/Lidocaine HCl 2% Spira-Wash™ Gel
- Misoprostol 0.0024%/Phenytoin 5%/Gentamicin 0.2% Spira-Wash™ Gel
- Tobramycin 5%/Mupirocin 5%/Itraconazole 2% Spira-Wash™ Gel
- Vancomycin 5%/Mupirocin 5% Spira-Wash™ Gel May be helpful with MRSA infections.
- Gentamicin 1 mg/Gm/Clindamycin 1 mg/Gm/Polymyxin 2,000 U/Gm Spira-Wash™ Gel
- Misoprostol 0.0024%/Phenytoin 5%/Metronidazole 2% Topical Gel
- Misoprostol 0.0024%/Phenytoin 5%/Gentamicin 0.2% Topical Gel

Treatment of Wounds with Pain

- Morphine Sulfate 1% Spira-Wash™ Gel
- Morphine Sulfate 1 mg/mL Topical Gel

Topical morphine is an option for wound pain and is usually requested after someone has had minimal success with lidocaine.

Additional Information to Consider

To help increase blood flow, consider nifedipine or pentoxifylline in or around the wound.

- Phenytoin 5%/Misoprostol 0.0024%/Nifedipine 2% Spira-Wash™ Gel
- Phenytoin 5%/Misoprostol 0.0024%/Nifedipine 2% Topical Gel (VersaBase®)

Spira-Wash™ Gel is a remarkably adherent water-washable wound care base. It is a soft opaque polyethylene glycol (PEG) ointment base containing organic meadowsweet extract. Due to the phenolic glycosides (spiraein) and flavonoids in the meadowsweet, it potentially has germicidal, anti-inflammatory and healing properties, thus making it a good choice for wounds, ulcers, burns, sores, and cuts. Since Spira-Wash is a water-washable base, it can easily be used in wounds that need to be cleaned/debrided on a regular basis.

Note: Spira-Wash should not be used on large areas of the body (~>20%).

Study results support the use of Spira-Wash Gel for wound management due to its high potential for inhibiting the growth of bacteria, with stronger action against Gram-negative bacteria. For more information, ask the pharmacist for the technical report.

