

PSORIASIS AND PSORIATIC ARTHRITIS SPECIALTY CARE PROGRAM

Phone: **856-963-4742** • Fax: **856-541-8580**



	NFORMATION:	PRESCRIBER INFORMATION:				
daress:	State: Zip: _	Address: State: Zip:				
	Alt. Phone:					
Παιι	Condor: O.M. O.E. Caragiyar:	NFI DEA				
oight:	Gender. O M O F Caregiver	Tax I.D.: Phone:				
eignt:	_ weight Allergies	Office Contact Priorie				
STATEMENT OF MEDICAL NECESSITY Date of Diagnosis:		atient also taking Methotrexate?				
טר-10: B Test: □ Positiv	/o D Nogative Date:	ep B ruled out or treatment started? Yes No Methotrexate				
FT: ALT:	AOT. D.1	pes patient have latex allergy?				
ssessment: Me		bes patient have joint involvement? Yes No Ses please indicate affected joint(s): Biologics				
% BSA affect	cted Li	3 //				
	J Chest ⊔ Arms ⊔ Hands ⊔ Nails │ _{fe}	f Prior Authorization is denied, recommended ormulary alternatives will be provided to the prescriber passed upon the patient's insurance coverage.				
NJECTION	TRAINING: O To be Administered by a Hea	althcare Provider O Pharmacist to Provide Training O Patient Trained in MD Office O Manufactor	urer Nurse	Support		
PICK UP O	R DELIVERY: O Delivery to Patie	nt's Home O Delivery to Physician's Office O Pharmacy to Coo	rdinate			
	N INFORMATION: (Please be sure	Front and Back Copies of Pharmacy and Medical Card to choose both induction and maintenance dose where applic Patient's Date of Birth:	able)			
Medication	Dosage & Strength	Direction	QTY	Refills		
		☐ Inject 400mg SC every other week				
⊒ CIMZIA [®]	☐ 200mg/ml Prefilled Syringe☐ 200mg Lyophilized Powder Vial	□ Inject 400 mg Sc every other week □ Induction Dose: (Weight ≤90kg) Inject 400 mg SC initially and at weeks 2 and 4 □ Maintenance Dose: (Weight ≤90kg) Inject 200 mg SC every other week Psoriatic Arthritis: □ Injitial: Inject 400 mg SC initially and at weeks 2 and 4 □ Maintenance: Inject 200 mg SC every other week □ Maintenance: Inject 400 mg SC every 4 weeks				
□ CIMZIA®		□ Induction Dose: (Weight ≤90kg) Inject 400mg SC initially and at weeks 2 and 4 □ Maintenance Dose: (Weight ≤90kg) Inject 200mg SC every other week Psoriatic Arthritis: □ Initial: Inject 400 mg SC initially and at weeks 2 and 4 □ Maintenance: Inject 200mg SC every other week	8	2		
	□ 200mg Lyophilized Powder Vial □ 50mg/ml Sureclick Autoinjector □ 50mg/ml Enbrel Mini™ Prefilled Cartridge For	□ Induction Dose: (Weight ≤90kg) Inject 400mg SC initially and at weeks 2 and 4 □ Maintenance Dose: (Weight ≤90kg) Inject 200mg SC every other week Psoriatic Arthritis: □ Initial: Inject 400 mg SC initially and at weeks 2 and 4 □ Maintenance: Inject 200mg SC every other week □ Maintenance: Inject 400mg SC every 4 weeks □ Induction Dose: Inject 50mg SC twice a week (3-4 days apart) for 3 months, then start maintenance dosing Maintenance: Inject 50mg SC once a week	8 4	2		
	□ 200mg Lyophilized Powder Vial □ 50mg/ml Sureclick Autoinjector □ 50mg/ml Enbrel Mini™ Prefilled Cartridge For Enbrel Mini™ only: AutoTouch™ Autoinjector	□ Induction Dose: (Weight ≤90kg) Inject 400mg SC initially and at weeks 2 and 4 □ Maintenance Dose: (Weight ≤90kg) Inject 200mg SC every other week Psoriatic Arthritis: □ Initial: Inject 400 mg SC initially and at weeks 2 and 4 □ Maintenance: Inject 200mg SC every other week □ Maintenance: Inject 400mg SC every 4 weeks □ Induction Dose: Inject 50mg SC twice a week (3-4 days apart) for 3 months, then start maintenance dosing □ Maintenance: Inject 50mg SC once a week Pediatric Patients: To achieve pediatric doses other than 50mg or 25mg, use reconstituted Enbrel lyophilized powder	4	2		
	□ 200mg Lyophilized Powder Vial □ 50mg/ml Sureclick Autoinjector □ 50mg/ml Enbrel Mini™ Prefilled Cartridge For	□ Induction Dose: (Weight ≤90kg) Inject 400mg SC initially and at weeks 2 and 4 □ Maintenance Dose: (Weight ≤90kg) Inject 200mg SC every other week □ Psoriatic Arthritis: □ Initial: Inject 400 mg SC initially and at weeks 2 and 4 □ Maintenance: Inject 200mg SC every other week □ Maintenance: Inject 400mg SC every 4 weeks □ Induction Dose: Inject 50mg SC twice a week (3-4 days apart) for 3 months, then start maintenance dosing □ Maintenance: Inject 50mg SC once a week □ Pediatric Patients: To achieve pediatric doses other than		2		
	□ 200mg Lyophilized Powder Vial □ 50mg/ml Sureclick Autoinjector □ 50mg/ml Enbrel Mini™ Prefilled Cartridge For Enbrel Mini™ only: AutoTouch™ Autoinjector □ 50mg/ml Prefilled Syringe □ 25mg/0.5ml Prefilled Syringe □ 25mg Lyophilized Powder Multiple Dose Vial □ Other:	□ Induction Dose: (Weight ≤90kg) Inject 400mg SC initially and at weeks 2 and 4 □ Maintenance Dose: (Weight ≤90kg) Inject 200mg SC every other week Psoriatic Arthritis: □ Initial: Inject 400 mg SC initially and at weeks 2 and 4 □ Maintenance: Inject 200mg SC every other week □ Maintenance: Inject 400mg SC every 4 weeks □ Induction Dose: Inject 50mg SC twice a week (3-4 days apart) for 3 months, then start maintenance dosing □ Maintenance: Inject 50mg SC once a week Pediatric Patients: To achieve pediatric doses other than 50mg or 25mg, use reconstituted Enbrel lyophilized powder □ > 138lbs or more: Inject 50mg weekly □ < 138lbs: Inject 0.8mg/kg weekly □ Other:	4	2		
	□ 200mg Lyophilized Powder Vial □ 50mg/ml Sureclick Autoinjector □ 50mg/ml Enbrel Mini™ Prefilled Cartridge For Enbrel Mini™ only: AutoTouch™ Autoinjector □ 50mg/ml Prefilled Syringe □ 25mg/0.5ml Prefilled Syringe □ 25mg Lyophilized Powder Multiple Dose Vial □ Other: □ Psoriasis 80mg/0.8ml and 40mg/0.4ml Starter Package	□ Induction Dose: (Weight ≤90kg) Inject 400mg SC initially and at weeks 2 and 4 □ Maintenance Dose: (Weight ≤90kg) Inject 200mg SC every other week Psoriatic Arthritis: □ Initial: Inject 400 mg SC initially and at weeks 2 and 4 □ Maintenance: Inject 200mg SC every other week □ Maintenance: Inject 400mg SC every 4 weeks □ Induction Dose: Inject 50mg SC twice a week (3-4 days apart) for 3 months, then start maintenance dosing □ Maintenance: Inject 50mg SC once a week Pediatric Patients: To achieve pediatric doses other than 50mg or 25mg, use reconstituted Enbrel lyophilized powder □ > 138lbs or more: Inject 50mg weekly □ < 138lbs: Inject 0.8mg/kg weekly	4 4			
	□ 200mg Lyophilized Powder Vial □ 50mg/ml Sureclick Autoinjector □ 50mg/ml Enbrel Mini™ Prefilled Cartridge For Enbrel Mini™ only: AutoTouch™ Autoinjector □ 50mg/ml Prefilled Syringe □ 25mg/0.5ml Prefilled Syringe □ 25mg Lyophilized Powder Multiple Dose Vial □ Other:	□ Induction Dose: (Weight ≤90kg) Inject 400mg SC initially and at weeks 2 and 4 □ Maintenance Dose: (Weight ≤90kg) Inject 200mg SC every other week Psoriatic Arthritis: □ Initial: Inject 400 mg SC initially and at weeks 2 and 4 □ Maintenance: Inject 200mg SC every other week □ Maintenance: Inject 400mg SC every 4 weeks □ Induction Dose: Inject 50mg SC twice a week (3-4 days apart) for 3 months, then start maintenance dosing □ Maintenance: Inject 50mg SC once a week □ Pediatric Patients: To achieve pediatric doses other than 50mg or 25mg, use reconstituted Enbrel lyophilized powder □ > 138lbs or more: Inject 50mg weekly □ < 138lbs: Inject 0.8mg/kg weekly □ Other: □ □ Induction Dose: Inject 80mg SC on day 1, then 40mg SC	4	0		
	□ 200mg Lyophilized Powder Vial □ 50mg/ml Sureclick Autoinjector □ 50mg/ml Enbrel Mini™ Prefilled Cartridge For Enbrel Mini™ only: AutoTouch™ Autoinjector □ 50mg/ml Prefilled Syringe □ 25mg/0.5ml Prefilled Syringe □ 25mg Lyophilized Powder Multiple Dose Vial □ Other: □ Psoriasis 80mg/0.8ml and 40mg/0.4ml Starter Package □ Psoriasis 40mg/0.4ml Starter Package □ 40mg/0.4ml Pen □ 40mg/0.4ml Prefilled Syringe □ Hidradenitis Suppurativa 80mg/0.8ml	□ Induction Dose: (Weight ≤90kg) Inject 400mg SC initially and at weeks 2 and 4 □ Maintenance Dose: (Weight ≤90kg) Inject 200mg SC every other week Psoriatic Arthritis: □ Initial: Inject 400 mg SC initially and at weeks 2 and 4 □ Maintenance: Inject 200mg SC every other week □ Maintenance: Inject 400mg SC every 4 weeks □ Induction Dose: Inject 50mg SC twice a week (3-4 days apart) for 3 months, then start maintenance dosing □ Maintenance: Inject 50mg SC once a week Pediatric Patients: To achieve pediatric doses other than 50mg or 25mg, use reconstituted Enbrel lyophilized powder □ > 138lbs or more: Inject 50mg weekly □ < 138lbs: Inject 0.8mg/kg weekly □ Other: □ Induction Dose: Inject 80mg SC on day 1, then 40mg SC on day 8, then 40mg SC every other week	4 4 3 4	0		
□ CIMZIA® □ ENBREL® □ HUMIRA®	□ 200mg Lyophilized Powder Vial □ 50mg/ml Sureclick Autoinjector □ 50mg/ml Enbrel Mini™ Prefilled Cartridge For Enbrel Mini™ only: AutoTouch™ Autoinjector □ 50mg/ml Prefilled Syringe □ 25mg/0.5ml Prefilled Syringe □ 25mg Lyophilized Powder Multiple Dose Vial □ Other: □ Psoriasis 80mg/0.8ml and 40mg/0.4ml Starter Package □ Psoriasis 40mg/0.4ml Starter Package □ 40mg/0.4ml Pen □ 40mg/0.4ml Prefilled Syringe □ Hidradenitis Suppurativa 80mg/0.8ml Starter Package □ Hidradenitis Suppurativa 40mg/0.4ml	□ Induction Dose: (Weight ≤90kg) Inject 400mg SC initially and at weeks 2 and 4 □ Maintenance Dose: (Weight ≤90kg) Inject 200mg SC every other week Psoriatic Arthritis: □ Initial: Inject 400 mg SC initially and at weeks 2 and 4 □ Maintenance: Inject 200mg SC every other week □ Maintenance: Inject 400mg SC every 4 weeks □ Induction Dose: Inject 50mg SC twice a week (3-4 days apart) for 3 months, then start maintenance dosing □ Maintenance: Inject 50mg SC once a week Pediatric Patients: To achieve pediatric doses other than 50mg or 25mg, use reconstituted Enbrel lyophilized powder □ > 138lbs or more: Inject 50mg weekly □ < 138lbs: Inject 0.8mg/kg weekly	4 3 4 2	0		
⊒ ENBREL®	□ 200mg Lyophilized Powder Vial □ 50mg/ml Sureclick Autoinjector □ 50mg/ml Enbrel Mini™ Prefilled Cartridge For Enbrel Mini™ only: AutoTouch™ Autoinjector □ 50mg/ml Prefilled Syringe □ 25mg/0.5ml Prefilled Syringe □ 25mg Lyophilized Powder Multiple Dose Vial □ Other: □ Psoriasis 80mg/0.8ml and 40mg/0.4ml Starter Package □ Psoriasis 40mg/0.4ml Starter Package □ 40mg/0.4ml Pen □ 40mg/0.4ml Prefilled Syringe □ Hidradenitis Suppurativa 80mg/0.8ml Starter Package □ Hidradenitis Suppurativa 40mg/0.4ml Starter Package	□ Induction Dose: (Weight ≤90kg) Inject 400mg SC initially and at weeks 2 and 4 □ Maintenance Dose: (Weight ≤90kg) Inject 200mg SC every other week Psoriatic Arthritis: □ Initial: Inject 400 mg SC initially and at weeks 2 and 4 □ Maintenance: Inject 200mg SC every other week □ Maintenance: Inject 400mg SC every 4 weeks □ Induction Dose: Inject 50mg SC twice a week (3-4 days apart) for 3 months, then start maintenance dosing □ Maintenance: Inject 50mg SC once a week Pediatric Patients: To achieve pediatric doses other than 50mg or 25mg, use reconstituted Enbrel lyophilized powder □ > 138lbs or more: Inject 50mg weekly □ < 138lbs: Inject 0.8mg/kg weekly □ Other: □ Induction Dose: Inject 80mg SC on day 1, then 40mg SC on day 8, then 40mg SC every other week □ Maintenance: Inject 40mg SC every other week □ Induction Dose: □ Inject two 80mg pens SC on day 1, then one 80mg pen SC on day 15 □ Inject one 80mg pen SC on day 1, one 80mg pen on day 2, then one 80mg pen on day 15	3 4 2 3 6	0 0		
⊒ ENBREL®	□ 200mg Lyophilized Powder Vial □ 50mg/ml Sureclick Autoinjector □ 50mg/ml Enbrel Mini™ Prefilled Cartridge For Enbrel Mini™ only: AutoTouch™ Autoinjector □ 50mg/ml Prefilled Syringe □ 25mg/0.5ml Prefilled Syringe □ 25mg Lyophilized Powder Multiple Dose Vial □ Other: □ Psoriasis 80mg/0.8ml and 40mg/0.4ml Starter Package □ Psoriasis 40mg/0.4ml Starter Package □ 40mg/0.4ml Pen □ 40mg/0.4ml Prefilled Syringe □ Hidradenitis Suppurativa 80mg/0.8ml Starter Package □ Hidradenitis Suppurativa 40mg/0.4ml	□ Induction Dose: (Weight ≤90kg) Inject 400mg SC initially and at weeks 2 and 4 □ Maintenance Dose: (Weight ≤90kg) Inject 200mg SC every other week Psoriatic Arthritis: □ Initial: Inject 400 mg SC initially and at weeks 2 and 4 □ Maintenance: Inject 200mg SC every other week □ Maintenance: Inject 400mg SC every 4 weeks □ Induction Dose: Inject 50mg SC twice a week (3-4 days apart) for 3 months, then start maintenance dosing □ Maintenance: Inject 50mg SC once a week Pediatric Patients: To achieve pediatric doses other than 50mg or 25mg, use reconstituted Enbrel lyophilized powder □ > 138lbs or more: Inject 50mg weekly □ < 138lbs: Inject 0.8mg/kg weekly □ Other: □ Induction Dose: Inject 80mg SC on day 1, then 40mg SC on day 8, then 40mg SC every other week □ Maintenance: Inject 40mg SC every other week □ Induction Dose: □ Induction Dose: □ Inject two 80mg pens SC on day 1, then one 80mg pen SC on day 15	3 4 2 3	0 0		
⊒ ENBREL®	□ 200mg Lyophilized Powder Vial □ 50mg/ml Sureclick Autoinjector □ 50mg/ml Enbrel Mini™ Prefilled Cartridge For Enbrel Mini™ only: AutoTouch™ Autoinjector □ 50mg/ml Prefilled Syringe □ 25mg/0.5ml Prefilled Syringe □ 25mg Lyophilized Powder Multiple Dose Vial □ Other: □ Psoriasis 80mg/0.8ml and 40mg/0.4ml Starter Package □ Psoriasis 40mg/0.4ml Starter Package □ 40mg/0.4ml Pen □ 40mg/0.4ml Prefilled Syringe □ Hidradenitis Suppurativa 80mg/0.8ml Starter Package □ Hidradenitis Suppurativa 40mg/0.4ml Starter Package □ Hidradenitis Suppurativa 40mg/0.4ml Starter Package □ 40mg/0.4ml Pen □ 40mg/0.4ml Prefilled Syringe	□ Induction Dose: (Weight ≤90kg) Inject 400mg SC initially and at weeks 2 and 4 □ Maintenance Dose: (Weight ≤90kg) Inject 200mg SC every other week Psoriatic Arthritis: □ Initial: Inject 400 mg SC initially and at weeks 2 and 4 □ Maintenance: Inject 200mg SC every other week □ Maintenance: Inject 400mg SC every 4 weeks □ Induction Dose: Inject 50mg SC twice a week (3-4 days apart) for 3 months, then start maintenance dosing □ Maintenance: Inject 50mg SC once a week Pediatric Patients: To achieve pediatric doses other than 50mg or 25mg, use reconstituted Enbrel lyophilized powder □ > 138lbs or more: Inject 50mg weekly □ < 138lbs: Inject 0.8mg/kg weekly	3 4 2 3 6	0 0		
⊇ ENBREL®	□ 200mg Lyophilized Powder Vial □ 50mg/ml Sureclick Autoinjector □ 50mg/ml Enbrel Mini™ Prefilled Cartridge For Enbrel Mini™ only: AutoTouch™ Autoinjector □ 50mg/ml Prefilled Syringe □ 25mg/0.5ml Prefilled Syringe □ 25mg Lyophilized Powder Multiple Dose Vial □ Other: □ Psoriasis 80mg/0.8ml and 40mg/0.4ml Starter Package □ Psoriasis 40mg/0.4ml Starter Package □ 40mg/0.4ml Pen □ 40mg/0.4ml Prefilled Syringe □ Hidradenitis Suppurativa 80mg/0.8ml Starter Package □ Hidradenitis Suppurativa 40mg/0.4ml Starter Package □ Hidradenitis Suppurativa 40mg/0.4ml Starter Package □ 40mg/0.4ml Pen □ 40mg/0.4ml Pen □ 40mg/0.4ml Prefilled Syringe	□ Induction Dose: (Weight ≤90kg) Inject 400mg SC initially and at weeks 2 and 4 □ Maintenance Dose: (Weight ≤90kg) Inject 200mg SC every other week Psoriatic Arthritis: □ Initial: Inject 400 mg SC initially and at weeks 2 and 4 □ Maintenance: Inject 200mg SC every other week □ Maintenance: Inject 400mg SC every 4 weeks □ Induction Dose: Inject 50mg SC twice a week (3-4 days apart) for 3 months, then start maintenance dosing □ Maintenance: Inject 50mg SC once a week Pediatric Patients: To achieve pediatric doses other than 50mg or 25mg, use reconstituted Enbrel lyophilized powder □ > 138lbs or more: Inject 50mg weekly □ < 138lbs: Inject 0.8mg/kg weekly □ Other: □ Induction Dose: Inject 80mg SC on day 1, then 40mg SC on day 8, then 40mg SC every other week □ Maintenance: Inject 40mg SC every other week □ Induction Dose: □ Inject two 80mg pens SC on day 1, then one 80mg pen SC on day 15 □ Inject one 80mg pen SC on day 1, one 80mg pen on day 2, then one 80mg pen on day 15 □ Maintenance: Inject 40mg SC on day 29 and every week thereafter □ Patient has signed HUMIRA Complete form	3 4 2 3 6	0 0		
□ ENBREL® □ HUMIRA® □ ORENCIA®	□ 200mg Lyophilized Powder Vial □ 50mg/ml Sureclick Autoinjector □ 50mg/ml Enbrel Mini™ Prefilled Cartridge For Enbrel Mini™ only: AutoTouch™ Autoinjector □ 50mg/ml Prefilled Syringe □ 25mg/0.5ml Prefilled Syringe □ 25mg Lyophilized Powder Multiple Dose Vial □ Other: □ Psoriasis 80mg/0.8ml and 40mg/0.4ml Starter Package □ Psoriasis 40mg/0.4ml Starter Package □ 40mg/0.4ml Pen □ 40mg/0.4ml Prefilled Syringe □ Hidradenitis Suppurativa 80mg/0.8ml Starter Package □ Hidradenitis Suppurativa 40mg/0.4ml Starter Package □ Hidradenitis Suppurativa 40mg/0.4ml Starter Package □ 40mg/0.4ml Pen □ 40mg/0.4ml Pen □ 40mg/0.4ml Prefilled Syringe All strengths	Induction Dose: (Weight ≤90kg) Inject 400mg SC initially and at weeks 2 and 4 Maintenance Dose: (Weight ≤90kg) Inject 200mg SC every other week Psoriatic Arthritis: Initial: Inject 400 mg SC initially and at weeks 2 and 4 Maintenance: Inject 200mg SC every other week Maintenance: Inject 200mg SC every 4 weeks Maintenance: Inject 400mg SC every 4 weeks Induction Dose: Inject 50mg SC twice a week (3-4 days apart) for 3 months, then start maintenance dosing Maintenance: Inject 50mg SC once a week Pediatric Patients: To achieve pediatric doses other than 50mg or 25mg, use reconstituted Enbrel lyophilized powder > 138lbs or more: Inject 50mg weekly < 138lbs: Inject 0.8mg/kg weekly Other: Induction Dose: Inject 80mg SC on day 1, then 40mg SC on day 8, then 40mg SC every other week Maintenance: Inject 40mg SC every other week Induction Dose: Inject 40mg SC every other week Inject two 80mg pens SC on day 1, then one 80mg pen SC on day 15 Inject one 80mg pen SC on day 1, one 80mg pen on day 2, then one 80mg pen on day 15 Maintenance: Inject 40mg SC on day 29 and every week thereafter Patient has signed HUMIRA Complete form Patient has signed HUMIRA Complete form Patient has signed HUMIRA Complete form Patient has signed Humira® Citrate Free	4 4 3 4 2 3 6	0 0		
⊒ ENBREL®	□ 200mg Lyophilized Powder Vial □ 50mg/ml Sureclick Autoinjector □ 50mg/ml Enbrel Mini™ Prefilled Cartridge For Enbrel Mini™ only: AutoTouch™ Autoinjector □ 50mg/ml Prefilled Syringe □ 25mg/0.5ml Prefilled Syringe □ 25mg Lyophilized Powder Multiple Dose Vial □ Other: □ Psoriasis 80mg/0.8ml and 40mg/0.4ml Starter Package □ Psoriasis 40mg/0.4ml Starter Package □ 40mg/0.4ml Pen □ 40mg/0.4ml Prefilled Syringe □ Hidradenitis Suppurativa 80mg/0.8ml Starter Package □ Hidradenitis Suppurativa 40mg/0.4ml Starter Package □ Hidradenitis Suppurativa 40mg/0.4ml Starter Package □ 40mg/0.4ml Pen □ 40mg/0.4ml Prefilled Syringe All strengths □ 125mg/ml ClickJect™ Autoinjector □ 125mg/ml Prefilled Syringe	Induction Dose: (Weight ≤90kg) Inject 400mg SC initially and at weeks 2 and 4 Maintenance Dose: (Weight ≤90kg) Inject 200mg SC every other week Psoriatic Arthritis: Initial: Inject 400 mg SC initially and at weeks 2 and 4 Maintenance: Inject 200mg SC every other week Maintenance: Inject 400mg SC every 4 weeks Induction Dose: Inject 50mg SC twice a week (3-4 days apart) for 3 months, then start maintenance dosing Maintenance: Inject 50mg SC once a week Pediatric Patients: To achieve pediatric doses other than 50mg or 25mg, use reconstituted Enbrel lyophilized powder > 138lbs or more: Inject 50mg weekly < 138lbs: Inject 0.8mg/kg weekly Other: Induction Dose: Inject 80mg SC on day 1, then 40mg SC on day 8, then 40mg SC every other week Maintenance: Inject 40mg SC every other week Other: Inject two 80mg pens SC on day 1, then one 80mg pen SC on day 15 Inject one 80mg pen SC on day 1, one 80mg pen on day 2, then one 80mg pen on day 15 Maintenance: Inject 40mg SC on day 29 and every week thereafter Patient has signed HUMIRA Complete form Sand dosages listed are Humira® Citrate Free Inject 125mg SC once a week Starter Pack: Take one tablet in the morning on day 1, then take one tablet in the	3 4 2 3 6 4	0 0 0		
DENBREL® HUMIRA®	□ 200mg Lyophilized Powder Vial □ 50mg/ml Sureclick Autoinjector □ 50mg/ml Enbrel Mini™ Prefilled Cartridge For Enbrel Mini™ only: AutoTouch™ Autoinjector □ 50mg/ml Prefilled Syringe □ 25mg/0.5ml Prefilled Syringe □ 25mg Lyophilized Powder Multiple Dose Vial □ Other: □ Psoriasis 80mg/0.8ml and 40mg/0.4ml Starter Package □ Psoriasis 40mg/0.4ml Starter Package □ 40mg/0.4ml Pen □ 40mg/0.4ml Prefilled Syringe □ Hidradenitis Suppurativa 80mg/0.8ml Starter Package □ Hidradenitis Suppurativa 40mg/0.4ml Starter Package □ 40mg/0.4ml Pen □ 40mg/0.4ml Pen □ 40mg/0.4ml Prefilled Syringe All strengths □ 125mg/ml ClickJect™ Autoinjector □ 125mg/ml Prefilled Syringe	Induction Dose: (Weight ≤90kg) Inject 400mg SC initially and at weeks 2 and 4 Maintenance Dose: (Weight ≤90kg) Inject 200mg SC every other week Psoriatic Arthritis: Initial: Inject 400 mg SC initially and at weeks 2 and 4 Maintenance: Inject 200mg SC every other week Maintenance: Inject 400mg SC every 4 weeks Induction Dose: Inject 50mg SC twice a week (3-4 days apart) for 3 months, then start maintenance dosing Maintenance: Inject 50mg SC once a week Pediatric Patients: To achieve pediatric doses other than 50mg or 25mg, use reconstituted Enbrel lyophilized powder > 138lbs or more: Inject 50mg weekly < 138lbs: Inject 0.8mg/kg weekly Other: Induction Dose: Inject 80mg SC on day 1, then 40mg SC on day 8, then 40mg SC every other week Maintenance: Inject 40mg SC every other week Other: Inject two 80mg pens SC on day 1, then one 80mg pen SC on day 15 Inject one 80mg pen SC on day 1, one 80mg pen on day 2, then one 80mg pen on day 15 Maintenance: Inject 40mg SC on day 29 and every week thereafter Patient has signed HUMIRA Complete form Starter Pack: Take one tablet in the morning on day 1, then take one tablet in the morning and one tablet in the evening as directed on the starter pack	3 4 2 3 6 4	0 0		



Signature:

Substitution Permitted

PSORIASIS AND PSORIATIC ARTHRITIS SPECIALTY CARE PROGRAM

Phone: **856-963-4742** • Fax: **856-541-8580**



Date:

Dispense As Written

PATIENT INFORMATION: Name:			PRESCRIBER INFORMATION: Name:				
City:	State: Zip	o:	Address: State: Zip:				
	Alt. Phone:				Fax:		
			NIDI.	DEA.			
OOB:	Gender: OM OF Caregiver	·	Tax I.D.:				
leight: V	Veight: Allergies:		Office Contact:	Phone:	Phone:		
3 STATEMENT	OF MEDICAL NECESSIT	Y: (Please Attach All I	Medical Documentation)	Prior Failed Treat			
Date of Diagnosis:	gnosis: Other: Serior		ent also taking Methotrexate?				
FB Test: ☐ Positive ☐ Negative Date: Does		Hep B ruled out o					
		Does patient have			☐ Oral Meds		
Assessment: Mode	erate Mod to Severe Severe	Does patient have	s patient have joint involvement? Yes No s, please indicate affected joint(s): Your Authorization is denied, recommended				
% BSA affected					☐ Biologics		
	Chest ☐ Arms ☐ Hands ☐ Nails		on is denied, recommended ves will be provided to the presc	riber			
, , , , , , , , , , , , , , , , , , ,			sed upon the patient's insurance coverage.		Others		
3 INJECTION T	RAINING: O To be Administered by a	Healthcare Provider O	Pharmacist to Provide Training O P	atient Trained in MD Office O Manu	facturer Nurs	e Support	
PICK UP OR	DELIVERY: O Delivery to Pa	atient's Home	Delivery to Physician's C	Office O Pharmacy to C	oordinat	е	
) INSURANCE	INFORMATION: Please In	clude Front and	Back Copies of Pharmac	y and Medical Card			
PRESCRIPTION Patient Name:	I INFORMATION: (Please b	oe sure to choos	e both induction and ma	aintenance dose where a	applicab	ole)	
Medication	Dosage & Strength		Direction		QTY	Refills	
☐ RASUVO®	Single-dose auto-injector prefilled syringe ☐ 7.5mg ☐ 10mg ☐ 12.5mg ☐ 15mg	e: 🔲 Inject	_ mg SC once weekly				
	☐ 17.5mg ☐ 10mg ☐ 12.5mg ☐ 15mg ☐ 27.5mg ☐ 20mg ☐ 22.5mg ☐ 25mg ☐ 27.5 mg ☐ 30mg		*An initial test dose of 2.5 to 5 mg is recommended in patients with risk factors for hematologic toxicity or renal impairment*				
SIMPONI® (for PsA)	□ 50mg/0.5ml Smartject Injector □ 50mg/0.5ml Prefilled Syringe	☐ Inject 50n	ng SC once a month		1		
		☐ Induction	☐ Induction Dose: Inject 150mg (two 75mg injections) SC at weeks 0 and 4		4	0	
☐ SKYRIZI [™]	☐ 75mg/0.83ml Prefilled Syringe	☐ Maintenan	☐ Maintenance: Inject 150mg (two 75mg injections) SC every 12 weeks thereafter				
	☐ Yes or ☐ No: SKYRIZI SELF-INJECTION: Healthcare provider certifies that patient has been trained and is eligible for self-injection						
☐ STELARA®	☐ 45 mg/0.5 mL Single-Dose Prefilled Sy☐ 45 mg/0.5 mL Solution in a Single-Dose	then every ringe Adult Dosi	asis: ng (≤100 kg): Inject 45 mg SC initial 12 weeks thereafter ng (>100 kg): Inject 90 mg SC initial 12 weeks thereafter				
	45 mg/6.5 mc 30lution in a Single-Dos	Psoriatic Art					
			ng SC initiation and at 4 weeks, then	n every 12 weeks thereafter			
☐ TREMFYA [™]	☐ 100mg/ml Prefilled Syringe		Dose: Inject 100mg/ml SC at we	eeks 0 and 4	2	0	
□ IKEWIFYA	☐ 100mg/ml One- Press Patient Controlle Injector		laintenance: Inject 100mg/ml SC every 8 weeks thereafter				
☐ XELJANZ [®]	□ 5mg Tablet	☐ Take one t	ablet by mouth twice daily in combi	nation with a nonbiologic DMARD	60		
☐ XELJANZ [®] XR	□ 11mg Tablet	☐ Take one t	ablet by mouth once daily in combi	nation with a nonbiologic DMARD	30		
<u> </u>							
	imzia®, Enbrel®, Humira®, Orencia™ aı	nd Otezla® are listed	alphabetically on respective en	rollment forms.	I		
	SIGNATURE: I authorize pharmacy		· · · · · · · · · · · · · · · · · · ·		t assistance n	rograms	

Prior authorization approval and insurance benefits will be determined by the payor based upon the patient's eligibility, medical necessity, and the terms of the patient's coverage, among other things. Participation in this program is not a guarantee of prior authorization or of payment

Signature: