

RHEUMATOID ARTHRITIS SPECIALTY CARE PROGRAM Phone: 856-963-4742 • Fax: 856-541-8580

KLOUDSCRIPT	
Community Led Specialty Pharmacy Care	ッ

O PATIENT INFORMATION:

PRESCRIBER INFORMATION:

			Name						
Address:			Address:						
	Stat				State: Zi	o:			
Phone:	Alt. Pho	ne:	Phone:		Fax:				
Email:			NPI:		_ DEA:				
	Gender: O M O F								
Height:	Weight: Al	lergies:	Office Contac	xt:	Phone:				
STATE	MENT OF MEDICAL NE	CESSITY: (Please Att	ach All Medical Documentation)		Prior Failed Treatmen				
Date of Diagnosis: Patient also taking Methotre				No Azulfidine [®] Biologics	Celebrex [®] Corticosteroids				

ICD-10:	Serious or active infection	on present?	🖵 Yes	🗆 No 🛛	Calcipotriene	🔲 Indocin [®]
	Hep B ruled out or treatr	ment started?	Yes	🗆 No	Indicate Drug	Name and Length of Treatment:
Other:	Does patient have latex	allergy?	Yes	🗆 No		
TB Test: Desitive Desitive Date:	LFT: ALT: AST	T: Da	ate:			

If Prior Authorization is denied, recommended formulary alternatives will be provided to the prescriber based upon the patient's insurance coverage.

4 INJECTION TRAINING: O Pharmacist to Provide Training O Patient Trained in MD Office O Manufacturer Nurse Support

PRODUCT DELIVERY: O Patient's Home O Physician's Office O Pharmacy to Coordinate

6 INSURANCE INFORMATION: Please Include Front and Back Copies of Pharmacy and Medical Card

PRESCRIPTION INFORMATION: (Please be sure to choose both induction and maintenance dose where applicable) Patient Name: Patient's Date of Birth:

Medication **Dosage & Strength** Direction QTY Refills <220 lbs: Inject 162mg SC every other week, followed by an increase to</p> □ 162mg/0.9ml Prefilled Syringe every week based on clinical response □ 162mg/0.9ml Prefilled Autoinjector (ACTPen[™]) □ ≥220 lbs: Inject 162mg SC every week □ Induction Dose: Inject 400mg SC on day 1, day 14 and day 28 6 0 Prefilled Syringe Starter Kit 200mg/ml Prefilled Syringe □ Maintenance: Inject 400mg SC every 4 weeks 200mg Lyophilized Powder Vial 2 □ Maintenance: Inject 200mg SC every other week □ 50mg/ml SureClick[®] Autoinjector 50mg/ml Enbrel Mini[®] Prefilled Cartridge For Enbrel Mini[®] only: AutoTouch[®] Autoinjector Inject 50mg SC once a week 50mg/ml Prefilled Syringe 25mg/0.5ml Prefilled Syringe 25mg/ml Vial 40mg/0.4ml Pen □ Inject 40mg SC every other week Patient has signed □ 40mg/0.4ml Prefilled Syringe □ Inject 40mg SC once a week HUMIRA Complete form All strengths and dosages listed are Humira[®] Citrate Free 150mg/1.14ml Prefilled Syringe 2 □ Inject 150mg SC every 2 weeks □ 150mg/1.14ml Prefilled Pen 200mg/1.14ml Prefilled Syringe □ Inject 200mg SC every 2 weeks 2 200mg/1.14ml Prefilled Pen Take one 2mg tablet by mouth with or without food daily 30 2mg Tablet Moderate Renal Impairment: Take one 1mg tablet by mouth with or □ 1mg Tablet 30 without food daily □ Induction Dose: <60 kg: 500mg administered IV, then inject 125mg SC within 24 hours</p> □ 125mg/ml Prefilled Syringe Go to 100 kg: 750mg administered IV, then inject 125mg SC within 24 hours 125mg/ml ClickJectTM Autoinjector □ >100 kg: 1000mg administered IV, then inject 125mg SC within 24 hours 250mg Lyophilized Powder Vial □ Inject 125mg SC once a week

Otezla®, Rasuvo®, Rinvoq™, Simponi®, Stelara®, Tremfya™, Xeljanz®, and Xeljanz®XR are listed alphabetically on respective enrollment forms.

	PRESCRIBER SIGNATURE: I authorize pharmacy to act as my designee for initiating and coordinating insurance prior authorizations, nursing services and patient assistance programs.						
	Signature:	Date:	Signature:	Date):		
	Substitution Permittee	1	5	Dispense As Written			
1	Prior authorization approval and insurance benefits will be determined by the p	payor based upon the patient's eligibility, medical necessity, and	the terms of the patient's coverage, and	ong other things. Participation in this program is not a guarantee of prior au	thorization or of payment.		

Confidentiality Notice: This fax is intended to be delivered only to the named addressee and contains confidential information that may be protected health information under federal and state laws. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please inform the sender immediately if you have received this document in error and then destroy this document immediately.



RHEUMATOID ARTHRITIS SPECIALTY CARE PROGRAM Phone: 856-963-4742 • Fax: 856-541-8580

KLOUDSCRIPT	
Community Led Specialty Pharmacy Care	ソ

PATIENT INFORMATION:

PRESCRIBER INFORMATION:

Name:			Na	me:			
Address:			Ad	dress:			
		_ State: Zip:	Cit	y:		_ State: Zip	o:
Phone:	A	t. Phone:	Pho	one:		Fax:	
Email:						DEA:	
		A OF Caregiver:					
Height:	Weight:	Allergies:	Off	ice Contact: _		Phone:	
	ENT OF MEDICA	L NECESSITY: (Please	se Attach All Medical Do	cumentation)	Pr	ior Failed Treatment	ts:
Date of Diagnosis:		Patient also takir	Patient also taking Methotrexate?		Azulfidine [®] Biologics	Celebrex [®] Corticosteroids	
ICD-10:		Serious or active	infection present?	🗆 Yes 🛛 No	Calcipotriene		

Hep B ruled out or treatment started?
Yes No Indicate Drug Name and Length of Treatment: Other: Does patient have latex allergy? □Yes □No TB Test: Desitive Negative Date: LFT: ALT: AST: Date:

If Prior Authorization is denied, recommended formulary alternatives will be provided to the prescriber based upon the patient's insurance coverage.

4 INJECTION TRAINING: O Pharmacist to Provide Training O Patient Trained in MD Office O Manufacturer Nurse Support

DRODUCT DELIVERY: O Patient's Home O Physician's Office O Pharmacy to Coordinate

(6) INSURANCE INFORMATION: Please Include Front and Back Copies of Pharmacy and Medical Card

PRESCRIPTION INFORMATION: (Please be sure to choose both induction and maintenance dose where applicable) Patient's Date of Birth:

Patient Name:

Dosage & Strength	Direction	QTY	Refills
Two-week Starter Pack (Titration)	□ Starter Pack: Take one tablet in the morning on day 1, then take one tablet in the morning and one tablet in the evening as directed on the starter pack	1	0
 28-day Starter Pack (Titration) 30mg tablets 	□ Maintenance: Take one 30mg tablet by mouth twice daily *For patients with severe renal impairment take one 30mg tablet once daily and skip afternoon doses in Starter Pack	60	
•	•		
□ 15mg Extended-Release Tablets	□ Take one 15mg tablet once a day	30	
 50mg/0.5ml Smartject® Autoinjector 50mg/0.5ml Prefilled Syringe 	□ Inject 50mg SC once a month	1	
	□ Induction Dose: Inject 45mg SC on day 1	1	
 45mg/0.5ml Prefilled Syringe 45mg/0.5ml Vial 90mg/1ml Prefilled Syringe 	□ Maintenance: Inject 45mg SC on day 29, and every 12 weeks thereafter	1	
	PsA with Coexistent Moderate-to-Severe Plaque Psoriasis (>220 lbs) Induction Dose: Inject 90 SC on day 1	1	
		1	
□ 100mg/ml Prefilled Syringe	□ Induction Dose: Inject 100mg/ml SC at weeks 0 and 4	2	0
100mg/ml One-Press Patient Controlled Injector	□ Maintenance: Inject 100mg/ml SC every 8 weeks thereafter	1	
	□ Take one 5mg tablet by mouth twice a day	60	
 5mg Tablet 11mg Tablet 	Take one 11mg tablet once a day *For patients with moderate renal or hepatic impairment take one 5mg tablet once daily	30	
	 Two-week Starter Pack (Titration) 28-day Starter Pack (Titration) 30mg tablets 30mg tablets 15mg Extended-Release Tablets 50mg/0.5ml Smartject® Autoinjector 50mg/0.5ml Prefilled Syringe 45mg/0.5ml Prefilled Syringe 45mg/0.5ml Vial 90mg/1ml Prefilled Syringe Yes or No: STELARA SELF-INJECTION: Healthcology 100mg/ml Prefilled Syringe 100mg/ml Prefilled Syringe 100mg/ml Prefilled Syringe 50mg/0.5ml Prefilled Syringe 	Image: Starter Pack (Titration) Starter Pack: Take one tablet in the morning on day 1, then take one tablet in the morning and one tablet in the evening as directed on the starter pack Image: Starter Pack (Titration) Maintenance: Take one 30mg tablet by mouth twice daily Somg tablets For patients with severe renal impairment take one 30mg tablet once daily and skip afternoon doses in Starter Pack Image: Starter Pack (Titration) Image: Starter Pack (Titration) Image: Starter Pack (Titration) Maintenance: Take one 30mg tablet by mouth twice daily Image: Starter Pack (Titration) Maintenance: Take one 30mg tablet by mouth twice daily Image: Starter Pack Image: Starter Pack Image: Starter Pack Take one 15mg tablet once a day Image: Starter Pack Image: Starter Pack Ima	Image: Starter Pack (Titration) Starter Pack: Take one tablet in the morning and one tablet in the evening as directed on the starter pack 1 28-day Starter Pack (Titration) Maintenance: Take one 30mg tablet by mouth twice daily 60 30mg tablets For patients with severe renal impairment take one 30mg tablet once daily and skip afternoon doses in Starter Pack 60 115mg Extended-Release Tablets Take one 15mg tablet once a day 30 650mg/0.5ml Smartject® Autoinjector Inject 50mg SC once a month 1 50mg/0.5ml Prefilled Syringe Induction Dose: Inject 45mg SC on day 1 1 45mg/0.5ml Vial Maintenance: Inject 45mg SC on day 29, and every 12 weeks thereafter 1 90mg/1ml Prefilled Syringe Induction Dose: Inject 90 SC on day 29, and every 12 weeks thereafter 1 100mg/ml Prefilled Syringe Induction Dose: Inject 90mg SC on day 29, and every 12 weeks thereafter 1 100mg/ml Prefilled Syringe Induction Dose: Inject 90mg SC on day 1 1 100mg/ml Prefilled Syringe Induction Dose: Inject 100mg/ml SC at weeks 0 and 4 2 100mg/ml Prefilled Syringe Induction Dose: Inject 100mg/ml SC at weeks 0 and 4 2 100mg/ml Prefilled Syringe Induction Dose: Inject 100mg/ml SC at weeks 0 and 4 2 100mg/ml Prefilled

Actemra®, Cimzia®, Colcigel®, Enbrel®, Humira®, Kevzara®, Olumiant® and Orencia® are listed alphabetically on respective enrollment forms.

PRESCRIBER SIGNATURE: I authorize pharmacy to act as my designee for initiating and coordinating insurance prior authorizations, nursing services and patient assistance programs.					
Signature:	Date:	Signature:	Date:		
Substitution Perm	litted	o	Dispense As Written		
Prior authorization approval and insurance benefits will be determined h	by the payor based upon the patient's eligibility, medical necessity, a	and the terms of the natient's coverage amo	ng other things. Participation in this program is not a guarantee of prior authorization or of payment		

Confidentiality Notice: This fax is intended to be delivered only to the named addressee and contains confidential information that may be protected health information under federal and state laws. If you are not the named addressee, you should not disseminate, distribute or copy this fax. Please inform the sender immediately if you have received this document in error and then destroy this document immediately.