



Est. 1976

Weight Loss

2704 N. Oak Street • Oak Center B-1 • Valdosta, GA 31602
Phone 229.244.5353 • Fax 229.244.5357
E-mail: hogansrx@gmail.com

Patient Name:	Prescriber:
Address:	Address:
City, State ZIP:	City, State ZIP:
Date of Birth:	Phone:
Phone #:	Fax:
Drug Allergies:	DEA #:

PRESCRIBER'S SIGNATURE: _____ DATE: _____

Semaglutide Sublingual

INITIAL TITRATION DOSE

- () **Semaglutide 1mg/ml Sublingual Suspension (QTY: 28ml)**
SIG: Place 0.3ml (0.3mg) under the tongue. Hold for 90 seconds then swallow once daily for 1 week, then increase to 0.5ml (0.5mg) daily thereafter. Do Not eat for 30 minutes

0.5mg MAINTENANCE DOSE

- () **Semaglutide 1mg/ml Sublingual Suspension (QTY: 28ml)**
SIG: Place 0.5ml under the tongue, hold for 90 seconds then swallow once daily.
Do Not eat for 30 minutes

1mg MAINTENANCE DOSE

- () **Semaglutide 1mg/ml Sublingual Suspension (QTY: 28ml)**
SIG: Place 1ml under the tongue, hold for 90 seconds then swallow once daily.
Do Not eat for 30 minutes

Refills _____

Q-Caps (Phentermine/Topiramate)

- () 6mg/23mg
- () 10mg/46mg
- () 14mg/69mg
- () 20mg/92mg

Sig: Take one capsule daily

Refills: _____