






Directions for Use:






1. Gather the materials you will need and place them on a table near the chair or bed where you will be sitting down: your medication vial, syringes, and alcohol swabs.
2. Wash hands with soap and water and then dry.
3. Open the alcohol swab and wipe the top of the vial. Allow the top to air dry.



4. Remove the syringe from its protective packaging.
5. Attach a needle if there is not already one.
6. While holding the syringe horizontally, pull the needle cover straight off and put it on the table.
7. Draw up an amount of air into the syringe equal to the amount of medication you will be drawing up. Insert the needle straight down into the rubber stopper of the vial and push in the air.



8. Turn the bottle and syringe upside down. Draw up the recommended dosage into the syringe. Tap the syringe gently and push the plunger to remove air bubbles.
9. Place the bottle and syringe right side up on the table. Gently remove the syringe from the vial and recap the needle.
10. Replace the needle with a fresh one.



11. You should be sitting upright or slightly reclined when performing the injection. You will be injecting this medication into the *corpora cavernosa*, the spongy tissue on the sides of the penis.
12. Choose an injection site on the side of the shaft of the penis (shaded area shown in the picture), in the first one-half area closest to the base of the penis. AVOID

VISIBLE BLOOD VESSELS.

13. Holding the head of your penis, stretch it lengthwise along your thigh so that you can clearly see the injection site
14. Clean the injection site with a new alcohol swab.
15. Remove the cover from the needle. Position the penis firmly against your thigh to keep it from moving during the injection.
16. Hold the syringe between your thumb and index finger. Using a steady motion, push the needle straight into the selected site until the metal part is almost entirely in the penis.
17. With a steady motion, push down on the plunger so that the entire volume of solution is slowly injected. Grasp the syringe barrel and pull the needle out of your penis.
18. Apply pressure to the injection site with the alcohol swab for about 5 minutes.
19. Place the used syringe in a sharps container for disposal. Do not reuse the syringe.

*****Remember: Alternate the side of the penis and vary the injection site with each use.**