

The Medication you have received contains Cholestyramine Resin Powder, Stevia (sweetener), Hydroxyethylcellulose (thickening agent), and Microcrystalline Cellulose (Filler)

- Keep this medication in the container it came in, tightly closed, and out of reach of children. Store it at room temperature and away from excess heat and moisture (not in the bathroom). Cholestyramine Powder should be discarded after 180 days.
- One level tablespoonful provides 4 grams of pure Cholestyramine Resin.
- Mix 1 level tablespoonful with 240ml (8oz) of water, apple juice or cranberry juice. Mix or blend thoroughly and let sit for 3 – 5 mins to allow it to thicken. (You may add more juice or water, but you need to drink the entire amount.)
- To prevent constipation, drink a full glass of water before and after you take the dose described above.
- Cholestyramine Powder should be taken 30 minutes prior to a meal. Take any other medication(s) either 1 hour before or 4 to 6 hours after taking Cholestyramine.
- This medication may cause constipation, bloating, gas, and/or abdominal discomfort. Please contact your physician regarding any concerns or issues that may occur after starting this medication.

The Medication you have received contains Cholestyramine Resin Powder, Stevia (sweetener), Hydroxyethylcellulose (thickening agent), and Microcrystalline Cellulose (Filler)

- Keep this medication in the container it came in, tightly closed, and out of reach of children. Store it at room temperature and away from excess heat and moisture (not in the bathroom). Cholestyramine Powder should be discarded after 180 days.
- One level tablespoonful provides 4 grams of pure Cholestyramine Resin.
- Mix 1 level tablespoonful with 240ml (8oz) of water, apple juice or cranberry juice. Mix or blend thoroughly and let sit for 3 – 5 mins to allow it to thicken. (You may add more juice or water, but you need to drink the entire amount.)
- To prevent constipation, drink a full glass of water before and after you take the dose described above.
- Cholestyramine Powder should be taken 30 minutes prior to a meal. Take any other medication(s) either 1 hour before or 4 to 6 hours after taking Cholestyramine.
- This medication may cause constipation, bloating, gas, and/or abdominal discomfort. Please contact your physician regarding any concerns or issues that may occur after starting this medication.

The Medication you have received contains Cholestyramine Resin Powder, Stevia (sweetener), Hydroxyethylcellulose (thickening agent), and Microcrystalline Cellulose (Filler)

- Keep this medication in the container it came in, tightly closed, and out of reach of children. Store it at room temperature and away from excess heat and moisture (not in the bathroom). Cholestyramine Powder should be discarded after 180 days.
- One level tablespoonful provides 4 grams of pure Cholestyramine Resin.
- Mix 1 level tablespoonful with 240ml (8oz) of water, apple juice or cranberry juice. Mix or blend thoroughly and let sit for 3 – 5 mins to allow it to thicken. (You may add more juice or water, but you need to drink the entire amount.)
- To prevent constipation, drink a full glass of water before and after you take the dose described above.
- Cholestyramine Powder should be taken 30 minutes prior to a meal. Take any other medication(s) either 1 hour before or 4 to 6 hours after taking Cholestyramine.
- This medication may cause constipation, bloating, gas, and/or abdominal discomfort. Please contact your physician regarding any concerns or issues that may occur after starting this medication.