


# Rectal Rocket Suppository Instructions

The Rectal Rocket was designed to be a comprehensive treatment for hemorrhoids. Part of the suppository is inserted into the rectum to treat internal hemorrhoids while the larger flared end of the suppository remains on the outside to treat external hemorrhoids.

- The suppository should be used when the patient will be able to lie down for at least six hours.
- The suppository should be at room temperature when placed in the rectum. Run the Rectal Rocket under warm water for a few seconds before insertion.
- Patients should consult their physician if they are experiencing continued excessive bleeding (bright red blood) from their hemorrhoids.

The diagram below identifies the proper placement of the rocket suppository in relationship to internal and external hemorrhoids.




# Rectal Rocket Suppository Instructions

The Rectal Rocket was designed to be a comprehensive treatment for hemorrhoids. Part of the suppository is inserted into the rectum to treat internal hemorrhoids while the larger flared end of the suppository remains on the outside to treat external hemorrhoids.

- The suppository should be used when the patient will be able to lie down for at least six hours.
- The suppository should be at room temperature when placed in the rectum. Run the Rectal Rocket under warm water for a few seconds before insertion.
- Patients should consult their physician if they are experiencing continued excessive bleeding (bright red blood) from their hemorrhoids.

The diagram below identifies the proper placement of the rocket suppository in relationship to internal and external hemorrhoids.



# Rectal Rocket Suppository Instructions

The Rectal Rocket was designed to be a comprehensive treatment for hemorrhoids. Part of the suppository is inserted into the rectum to treat internal hemorrhoids while the larger flared end of the suppository remains on the outside to treat external hemorrhoids.

- The suppository should be used when the patient will be able to lie down for at least six hours.
- The suppository should be at room temperature when placed in the rectum. Run the Rectal Rocket under warm water for a few seconds before insertion.
- Patients should consult their physician if they are experiencing continued excessive bleeding (bright red blood) from their hemorrhoids.

The diagram below identifies the proper placement of the rocket suppository in relationship to internal and external hemorrhoids.

