Bell Pharmacy

Myasthenia Gravis Enrollment Form (gMG)

1907 State Route 27, Edison, NJ 08817 | www.bellpharmacy.net | Email: contact@bellpharmacy.net | Phone: 732-985-1211 | Fax: 732-985-3609

PATIENT INFORMATION

Name:	Date of Birth:		O Male O Female
Address:	City:	State:	Zip:
Phone:	Alt Phone:		Email:
SS #:	Primary Language:		Emergency Contact:

PRESCRIBER INFORMATION

Prescribing Practitioner:				NPI#:
Supervising Physician:				NPI#:
Address:	City:	State:	Zip:	Tax ID:
Phone:	Fax:			Office Contact:

PRESCRIPTION INFORMATION

Needs by Date: Ship to: O Patients home O Prescriber 1st order only O Prescriber all orders O Other

Drug	Dose	Direction & Quantities	Refills
O Intravenous Immunoglobulin	O 0.4 gm/kg O 1gm/kg O 2gm/kg O ________ grams	Infuse: O IV daily x ____ day(s); repeat every ______ week(s) x ______ cycles O Other: ____	
O Soliris	O 300 mg/30 mL vial (10 mg/mL)	O Dose Titration – Month 1: Administer 900 mg via IV infusion every 7 days for 4 weeks. 4 week-supply	0
		O Maintenance Dosing: Administer 1,200 mg via IV infusion every 2 weeks starting Week 5 O 4-week supply O 12-week supply O Other___________	1 - year supply
O Pre-medications:	O Acetaminophen 650mg PO 30 mins prior to infusion		
	O Diphenhydramine 25mg PO 30 mins prior to infusion		

O Other:

O Other Pre-medications:

MEDICAL INFORMATION

** PLEASE FAX COPY OF PRESCRIPTION MEDICATION/MEDICAL CARD, FRONT AND BACK, AS WELL AS ANY CLINICAL NOTES REGARDING THERAPY **

Patient Height:____________________ in/cm Weight: ____________________ kg/lbs Allergies: _____________________ Bun/Creat.:____________________

Diagnosis	ICD-10
Neuromuscular:	
O Myasthenia Gravis with (Acute) Exacerbation	G70.01
O Myasthenia Gravis (MG)	G70.0

Please Draw: O CBC/diff O CMP O IgG w/ subclasses 1-4 O Quant. Ig O ___________ O __________ Frequency:__________

PER Anaphylaxis Protocol:
O Adult – EpiPen 0.3 auto-injector dual pack
O Pediatric - EpiPen 0.15 auto-injector dual pack

Administer intramuscularly in the event of ADR
[May repeat x 1. Order is valid for 1 year]. **Use generic if applicable**

If applicable, flush intravenous access device per Parkway Pharmacy protocol:

Access	NS	Heparin 100 u/ml
Peripheral	1-3ml before/after use	10u/ml 1-2mls after last NS flush
Midline, central (non-port), PICC	NS 5-10 mls before/after use; 10mls after blood draw	10 u/ml 3-5mls after last NS flush; 5mls after blood draw
Implanted Port	5-10mls before/after use; 20mls after blood draw	100 u/ml 5mls after last NS flush; 5mls after blood draw
Tunneled	5-10mls before/after use; 20mls after blood draw	10 u/ml 3- mls after last NS flush. 5mls after blood draw
Groshong PICC, Midline	5-10mls before/after use; 10mls after blood draw	NO Heparin needed

PRESCRIBING PRACTITIONER SIGNATURE

To Prescribing Practitioner: By signing this form and utilizing our services, you are also authorizing Parkway Pharmacy to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies, and co-pay assistance foundations.

Prescribing Practitioner: Date

CONFIDENTIALITY NOTICE

IMPORTANT: This fax is intended to be delivered only to the named addressee. It contains material that is confidential, proprietary or exempt from disclosure under applicable law. If you are not the named addressee, you should not disseminate, distribute, or copy this fax. Please notify the sender immediately if you have received this document in error and then destroy this document immediately.

Faxed Prescriptions will only be accepted from a prescribing practitioner.