

# Myristin<sup>®</sup> Equine Hip & Joint Supplement

## Recommended Feeding Regimen

Myristin<sup>®</sup> Equine Hip & Joint Supplement powder is sold in 1 kilogram bags.  
 A level measuring tablespoonful of powder contains 12.5 grams, of which 2.5 grams are molecular Cetyl Myristoleate (CMO).

Daily feeding instructions		
<p><b>First 60 days</b></p> <p>By the end of the 60 days, your horse will have taken the recommended initial regimen of 600 grams of molecular Cetyl Myristoleate (CMO)</p>	<p>Give 2 measuring tablespoonfuls, mixed with sweet feed or similar food, in first feeding of the day.</p> <p>Give same amount with another feeding, later in the day.</p> <p>(provides a total of 10 grams of molecular CMO per day)</p>	<p><u>How much should I order?</u>            Initial 60-day regimen requires a total of 3 bags (1 kg each) of Myristin<sup>®</sup> Equine Hip &amp; Joint Supplement</p>
<p><b>Maintenance Regimen</b></p> <p>This maintenance regimen offers your horse an approximate monthly dose of 150 grams of molecular Cetyl Myristoleate (CMO)</p>	<p>Give 1 measuring tablespoonful, mixed with sweet feed or similar food, in first feeding of the day.</p> <p>Give same amount with another feeding, later in the day.</p> <p>(provides a total of 5 grams of molecular CMO per day)</p>	<p><u>How much should I order?</u>            Each month of the maintenance regimen requires 3/4 of a bag of Myristin<sup>®</sup> Equine Hip &amp; Joint Supplement</p>

### Feeding Tips:

Mix Myristin<sup>®</sup> Equine Hip & Joint Supplement powder with any kind of sweet feed, oats, or similar food. If horse is reluctant to eat the powder, starting with a smaller dose per feeding will help familiarize horse with powder. If horse will not eat powder when mixed with sweet feed, try moistening the feed so that powder adheres to it. If horse still rejects powder, try mixing with molasses, apple sauce, or wet beet pulp. For difficult eaters, syringe in the first few feedings mixed with applesauce until the horse accepts product.

### Important:

**Keep out of reach of children. In case of accidental overdose in a human or animal, contact the appropriate healthcare professional. Consider consulting with your veterinarian before giving to breeding or pregnant horses.**

*This statement has not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.*