

Myristin® Hip & Joint Formula for Pets

Feeding Instructions

Dogs



Loading Dose Amount:

Give one tablet per day per 25 pounds of weight for two months (see chart below). Older animals may need to be given the loading amount for longer than two months.

Recommended number of tablets to give based on dog's weight	
up to 25 lbs	1 tablet daily
25-50 lbs	2 tablets daily
51-75 lbs	3 tablets daily
76-99 lbs	4 tablets daily
100-149 lbs	5 tablets daily
150 lbs and over	6 tablets daily


Best given at the time of a regular daily feeding. Tablets can be given half in the AM and half in the PM. Tablets may be crushed for very small animals. If desired, the tablets may be given as a treat on an empty stomach an hour before feeding with water available.

Maintenance Dose Amount:

Recommended maintenance amounts are one-half of the loading amounts:

Recommended number of tablets to give based on dog's weight	
Up to 25 lbs	1/2 tablet daily
25-50 lbs	1 tablet daily
51-75 lbs	1 1/2 tablets daily
76-99 lbs	2 tablets daily
100-149 lbs	2 1/2 tablets daily
150 lbs and over	3 tablets daily

Cats



Loading Dose Amount:

Give 1 tablet daily for 60 days. Tablets may need to be crushed and given on food.

Maintenance Dose Amount:

Give 1/2 tablet daily.

Consider consulting with your veterinarian before giving to breeding or pregnant animals. If tablets are not eaten readily, crush and add to food.

Important: Keep out of reach of children. In case of accidental overdose in a human or animal, contact the appropriate healthcare professional.